

A Study on Fuzzy Travelling Salesman Problem Using Fuzzy Number

Dr. Amit Kumar Rana

Assistant Professor, Department of Mathematics, Keral Verma Subharti College of Science,
Swami Vivekanand Subharti University, Meerut, Uttar Pradesh, INDIA.

rana2977@gmail.com

ABSTRACT - A fuzzy number can be used to solve many real-life problems like Travelling Salesman, Assignment Problems and so on. Travelling Salesman Problem (TSP) solutions are based on the assumption that time of travel between nodes is wholly dependent on distance. But in practice, this is not so as road and traffic conditions help to determine time taken to travel between nodes. We introduce fuzzy based TSP solution where distance and road traffic conditions are fuzzy. In this paper, a new method is proposed for solving Travelling Salesman problems using Transitive Fuzzy Numbers. We use Haar Hungarian algorithm approach to solve a Fuzzy Travelling Salesman Problem (FTSP) and Numerical examples are given to validate the proposed algorithm.

KEYWORDS: Fuzzy numbers, Transitive Trapezoidal Fuzzy Numbers, Haar Ranking, Hungarian Method, Modeling, Optimization.

I. INTRODUCTION

Traveling salesman problem is a classical problem in combinatorial optimization. The objective of the problem is to find the shortest route of salesman starting from a given city, visiting all other cities only once and finally come to the same city where he started. There are different approaches for solving traveling salesman problems. Linear programming method, heuristic methods like cutting plan algorithms and branch and bound method, Markov chain, simulated annealing and so on.

In real life situation, it may not be possible to get the cost or time as a certain quantity. To overcome this [1] introduce fuzzy set concepts to deal with imprecision and vagueness. Since then significant advantages have been made in developing numerous methodologies and their applications to various decision problems. If the cost or time or distance is not crisp values, then it becomes a fuzzy TSP. The Fuzzy TSP has been solved for LR-fuzzy parameters by [2]. In this paper, a new algorithm which is similar to the classical assignment method is introduced to solve fuzzy TSP. First, Fuzzy Hungarian method is applied and then modifications are done to satisfy the route conditions by considering element-wise subtraction and [3] ranking method of fuzzy numbers. [4] solved fuzzy TSP by using the Hungarian algorithm with element-wise subtraction of fuzzy numbers. Since fuzzy TSP is an NP-hard problem, more numbers of algorithms are still proposed to solve the fuzzy TSP. In this paper, the ordering of fuzzy numbers based on Haar wavelet is used [5], to order the fuzzy numbers. The advantage of using this ranking technique is that it converts a given fuzzy number into average and

detailed coefficients using down sampling. The uniqueness of the Haar ranking method [5] is that the fuzzification from the defuzzified value is very easy to obtain through up sampling. In this paper, the Hungarian method using Haar tuples [6] is proposed for solving traveling salesman problem with fuzzy parameters. A New Approach is used by [7] to Solve Fuzzy Travelling Salesman Problems by using ranking functions. A study on travelling salesman problems using dynamic programming algorithm studied by [8]. Fuzzy multi objective linear programming approach used by [9] for solving travelling salesman problem. [10] studied application of fuzzy ranking method for solving assignment problems with fuzzy costs. A study Travelling salesman problem by using a fuzzy multi-objective linear programming was carried out by [11]. Thus fuzzy number and fuzzy linear programming is a useful tool for solving travelling salesman problems.

II. PRELIMINARIES

Definition 2.1

Fuzzy sets unlike crisp sets are characterized by degree of membership. For example in a crisp set of tall people, one can either belong to it or not. But in a fuzzy set of tall people the membership is characterized by values which range from 1 for complete membership to zero for non membership. Hence, giants may belong to this to a degree of one while dwarfs belong to this to a degree of zero

A notation convention for fuzzy sets when the universe of discourse, X , is discrete and finite, is as follows for a fuzzy set

$$A = \frac{\mu_A(x_1)}{x_1} + \frac{\mu_A(x_2)}{x_2} + \dots$$

$$= \frac{\mu_A(x_i)}{x_i}$$

Definition 2.2

A Fuzzy number \hat{A} with membership function satisfies piece wise continuity, convexity and normality.

Definition 2.3 A fuzzy number $\hat{A} = (a, b, c, d,)$ with membership function defined as

$$\mu_n(x) = \left\{ \begin{array}{ll} \left(\frac{(x-a)}{(b-a)} \right) & a \leq x \leq b \\ 1 & b \leq x \leq c \\ \left(\frac{(d-x)}{(d-c)} \right) & c \leq x \leq d \\ 0 & otherwise \end{array} \right\}$$

is known as trapezoidal fuzzy number.

Definition 2.4:

The Fuzzy Operations of fuzzy numbers are defined as

Fuzzy Addition:

$$(a_1, b_1, c_1, d_1) + (a_2, b_2, c_2, d_2) = (a_1 + a_2, b_1 + b_2, c_1 + c_2, d_1 + d_2)$$

Fuzzy Subtraction:

$$(a_1, b_1, c_1, d_1) - (a_2, b_2, c_2, d_2) = (a_1 - a_2, b_1 - b_2, c_1 - c_2, d_1 - d_2)$$

Definition 2.5:

If $a(1) \sim a(4)$, then the trapezoidal fuzzy number $A = (a(1), a(2), a(3), a(4))$ is called transitive trapezoidal fuzzy number. It is denoted by $A = (a(1), a(4))$, Where $a(1)$ is core (A), $a(4)$ is left width and right width of c.

The parametric form of a transitive trapezoidal fuzzy number is represented by

$$A = [a(1) - a(2)(1-r), a(1)+a(3)(1-r), a(1) - a(4)(1-r)]$$

Definition 2.6:

Consider a Trapezoidal fuzzy number $\hat{A} = (a, b, c, d,)$, the Haar tuple of \hat{A} can be calculated as $H(\hat{A}) = (\alpha, \beta, \gamma, \delta)$ where $\alpha = (a+b+c+d)/4$, $\beta = [(a+b)-(c+d)]/4$, $\gamma = (a-b)/2$, $\delta = (c-d)/2$. Here 'a' is the average coefficient and the remaining coefficients are all called as detailed coefficients of the given trapezoidal fuzzy number.

Definition 2.7:

Element wise Addition:

$$(\alpha_1, \beta_1, \gamma_1, \delta_1) + (\alpha_2, \beta_2, \gamma_2, \delta_2) = (\alpha_1 + \alpha_2, \beta_1 + \beta_2, \gamma_1 + \gamma_2, \delta_1 + \delta_2)$$

Element wise Subtraction:

$$(\alpha_1, \beta_1, \gamma_1, \delta_1) - (\alpha_2, \beta_2, \gamma_2, \delta_2) = (\alpha_1 - \alpha_2, \beta_1 - \beta_2, \gamma_1 - \gamma_2, \delta_1 - \delta_2)$$

III. FUZZY TRAVELLING SALESMAN PROBLEM

Suppose a person has to visit n cities. He starts from a particular city, visits each city once and then returns to the starting point. The fuzzy travelling costs from i th city to j th city is given by C_{ij} . The chosen fuzzy travelling salesman problem may be formulated as

$$\text{Minimize } Z = \sum_{i=0}^n \sum_{j=0}^n c_{ij} x_{ij}$$

subject to

$$\sum_{i=0}^n x_{ij} = 1 \text{ and } \sum_{j=0}^n x_{ij} = 1$$

3.1 Proposed Algorithm

Consider the matrix form of the fuzzy travelling salesman problem.

Step: 1 Transform the given fuzzy parameters into Haar tuples.

Step: 2 Fuzzy optimum assignment is obtained by using the Hungarian algorithm.

Step: 3 Inspecting the obtained solution to see whether the solution satisfied route conditions or not. If it is then it is the optimal solution of fuzzy TSP. If not, take the next best solution.

3.2 Numerical Example

Consider the following fuzzy TSP discussed in [7] is given in table 1

Table 1

CITY	A	B	C	D
A	∞	(4,3,3,4)	(1,4,2,1)	(5,13,1,5)
B	(2,6,3,2)	∞	(6,4,2,6)	(2,3,7,2)
C	(5,13,1,5)	(5,6,7,5)	∞	(1,4,2,1)
D	(6,12,1,6)	(1,4,2,1)	(2,6,3,2)	∞

Convert the given fuzzy numbers into Haar tuples. The TSP with Haar tuples as its elements is given in the table 2

Table 2

CITY	A	B	C	D
A	∞	(3.5,0,0.5, -0.5)	(2, 0.5, -1.5,0.5)	(6, 3,-4,-2)
B	(3.25,0.75,-2,0.5)	∞	(4.5,0.5,1,-2)	(3.5,-1,-.5,2.5)
C	(6,3,-4,-2)	(5.75,-.25,-0.5,1)	∞	(2,0.5,-1.5,0.5)
D	(6.25,2.75,-3,-2.5)	(2, 0.5, -1.5,0.5)	(3.25,1.5,-2,0.5)	(2,0.5,-1.5,0.5)

now apply the Hungarian algorithm to get the table 3 as

Table 3

CITY	A	B	C	D
A	∞	(1.5,0.5,2,-1)	(0,0,0,0)	(4, 2.5,-2.5,-2.5)
B	(0,0,0,0)	∞	(1.25,-.25,3,-.25)	(.25,-1.75,1.5,2)
C	(4,2.5, -2.5, -2.5)	(3.75, 0.75,1,.5)	∞	(0,0,0,0)
D	(4.25,2.25, -1.5,-3)	(0,0,0,0)	(1.25,1.5,-.5,0)	∞

The optimal assignment schedule is A→C→D→B→A. Here the route condition is satisfied.

Haar optimal cost is $(2,0.5,-1.5,0.5)+(2,0.5,-1.5,0.5)+(2,0.5,-1.5,0.5)+(3.25,0.75,-2,0.5) = (9.25,2.25,-6.5,2)$

and the corresponding fuzzy optimal cost after defuzzification is given by $(5,18,9,5) = \text{Rs.}9.28$

IV. CONCLUSION

An Algorithm using Haar tuples is proposed for solving transitive fuzzy traveling salesman problem. Since this algorithm is similar to the Hungarian algorithm it is effective and easy to compute. This proposed method is effective than the existing methods because the results were obtained in Haar parameters and further it can be transformed into parameters. The results match with the existing technique and also satisfy the regular TSP conditions and thus provide a platform for further study on TSP.

REFERENCES

[1] RE Bellman, LA Zadeh .. “Decision-making in a fuzzy environment,” *Manage. Sci.*, Vol.17: pp141-164. 1970.

[2] Amit kumar ,Anil Gupta “Assignment and travelling salesman problems with Coefficient as LR parameters, ” *Int. Journal of Applied Science and Engineering*, Vol.10, No.3,pp155-170. 2012

[3] R.R.Yager, “A procedure for ordering fuzzy subsets of the unit interval,” *Information Sciences*, 24,pp 143-161. 1981

[4] S. Dhanasekar, S. Hariharan and P. Sekar . “Classical Travelling Salesman problem (TSP) based approach to solve fuzzy TSP using Yager’s ranking,” *International Journal of Computer Applications (IJCA)*, Vol 74(13),pp1-4. 2013.

[5] S. Dhanasekar, S. Hariharan and P. Sekar. “Ranking of Generalized Trapezoidal fuzzy numbers using Haar wavelet “, *Applied Mathematical Sciences*, Vol.8, No.160,pp7951-7958, 2014.

[6] S. Dhanasekar, S. Hariharan and P. Sekar. “Haar Hungarian algorithm to solve fuzzy Assignment problem” *International Journal of Pure and Applied Mathematics*, Vol.113, No.7, pp58-66, 2017.

[7] S. Chandrasekaran, G. Kokila, Junu Saju , “ A New Approach to Solve Fuzzy Travelling Salesman Problems by using ranking functions “, *International Journal of Science and Research*, Vol.10, No.3,pp155-170. 2012.

[8] A. Singhal, P. Pandey. “Travelling Salesman Problems by Dynamic Programming algorithm,” *International Journal of Scientific Engineering and Applied Science*, Vol.2,pp263 – 267, 2016.

[9] A. Rehmat, H. Saeed, M S Cheema “Fuzzy multi - objective linear programming approach for travelling salesman problem” *Pakistan Journal of Statistics*, Vol.3, No.2,pp87-98, 2007.

[10] S. Mukherjee, K. Basu.”Application of fuzzy ranking method for solving assignment problems with fuzzy costs”. *International Journal of Computational and Applied Mathematics*, Vol.5:pp 359-368.2010

[11] Sepideh Fereidouni. “Travelling salesman problem by using a fuzzy multi-objective linear programming”. *African Journal of mathematics and computer science research* ,Vol.4(11),pp 339-349,2011.